



IMMUNIZATIONS/SITE: 12-35 Months (Toddlers)

Recommended IM Site: Vastus Lateralis (thigh)

SUBQ Site: Back of arm

VACCINE: LEFT	ROUTE	VACCINE: RIGHT
DTaP/DT Pediarix (DTaP+IPV+HepB) Pentacel (DTaP+IPV+Hib) *Use anterolateral thigh	IM	
IPV (Polio)	SUBQ/IM (arm)	
VARICELLA	SUBQ (arm)	
 *PPSV23 can be given Sub Q or IM	IM	HEPATITIS A If receiving >2 IMs in same site, can move Left
	IM	INACTIVATED INFLUENZA (0.25mL) (Flu also comes in intranasal for 2yrs+)
	IM	Hib Comvax (Hib+HepB) MenHibrix (mening. +hib) If receiving >2 IMs in same site, can move Left
	IM	HEPATITIS B
	IM	PCV 13 (Pneumococcal Conjugate) *Use anterolateral thigh
	SUBQ (arm)	MMR
	SUBQ (arm)	PROQUAD (MMR+VAR) (through age 12)

*Tetanus-containing vaccines and PCV-13 are most irritating to tissues and should be in separate limbs- preferably thighs
 -Multiple vaccines are better handled in the vastus lateralis due to the greater muscle mass, though may use deltoid for single IM if muscle mass is adequate.

