



IMMUNIZATIONS/SITE: 3yrs+ (Child-Adult)

IM Site: Deltoid

SUBQ Site: Back of arm

VACCINE: LEFT	ROUTE	VACCINE: RIGHT
**Tdap/Td (7yrs+ only) DTaP/DT (under 7yrs only) Pediarix (DTaP+HepB+IPV) (thru 6yrs) Pentacel (DTaP+IPV+Hib) (thru 4yrs) Kinrix (DTaP+IPV) (4-6yrs)	IM	
IPV (Polio)	SUBQ	
VARICELLA	SUBQ	
 *PPSV23 can be given Sub Q or IM	IM	HEPATITIS A HEPATITIS B If receiving >2 IMs in one limb, can move to Left
	IM	**MCV4 (Meningococcal) * Give Men B in left
	IM	Inactivated Influenza (0.5mL) (Flu also comes in intranasal for 2yrs+)
	IM	Hib If receiving >2 IMs in one limb, can move to Left
	IM	PCV 13 (Pneumococcal Conjugate) PPVS23 (Pneumococcal Polysaccharide)
	IM	**HPV (Human Papillomavirus)
	SUBQ	MMR
	SUBQ	PROQUAD (MMR+VAR) (through age 12)

****When giving Tdap, MCV4, and HPV, give most reactive (Tdap and MCV4) in separate limbs and give HPV last (most painful).*** If giving MCV4 & Men B give in separate arms. Adolescents are more prone to syncope. Best practice is to have them remain seated x15-20 minutes to ensure safety.

