

Administering Vaccines:

Dose, Route, Site, and Needle Size


Vaccine	Dose	Route
Diphtheria, Tetanus, Pertussis (DTaP, DT, Tdap, Td)	0.5 mL	IM
<i>Haemophilus influenzae</i> type b (Hib)	0.5 mL	IM
Hepatitis A (HepA)	≤18 yrs: 0.5 mL ≥19 yrs: 1.0 mL	IM
Hepatitis B (HepB) Persons 11–15 yrs may be given Recombivax HB (Merck) 1.0 mL adult formulation on a 2-dose schedule.	≤19 yrs: 0.5 mL ≥20 yrs: 1.0 mL	IM
Human papillomavirus (HPV)	0.5 mL	IM
Influenza, live attenuated (LAIV)	0.2 mL (0.1 mL in each nostril)	Intranasal spray
Influenza, inactivated (IIV); recombinant (RIV), for ages 18 years and older	6–35 mos: 0.25 mL ≥3 yrs: 0.5 mL	IM
Influenza (IIV) Fluzone Intradermal, for ages 18 through 64 years	0.1 mL	ID
Measles, Mumps, Rubella (MMR)	0.5 mL	Subcut
Meningococcal conjugate (MCV4 [MenACWY])	0.5 mL	IM
Meningococcal serogroup B (MenB)	0.5 mL	IM
Meningococcal polysaccharide (MPSV)	0.5 mL	Subcut
Pneumococcal conjugate (PCV)	0.5 mL	IM
Pneumococcal polysaccharide (PPSV)	0.5 mL	IM or Subcut
Polio, inactivated (IPV)	0.5 mL	IM or Subcut
Rotavirus (RV)	Rotarix: 1.0 mL Rotateq: 2.0 mL	Oral
Varicella (Var)	0.5 mL	Subcut
Zoster (Zos)	0.65 mL	Subcut
Combination Vaccines		
DTaP-HepB-IPV (Pediarix) DTaP-IPV/Hib (Pentacel) DTaP-IPV (Kinrix; Quadracel) Hib-HepB (Comvax) Hib-MenCY (MenHibrix)	0.5 mL	IM
MMRV (ProQuad)	≤12 yrs: 0.5 mL	Subcut
HepA-HepB (Twinrix)	≥18 yrs: 1.0 mL	IM

Injection Site and Needle Size		
Subcutaneous (Subcut) injection Use a 23–25 gauge needle. Choose the injection site that is appropriate to the person's age and body mass.		
AGE	NEEDLE LENGTH	INJECTION SITE
Infants (1–12 mos)	5/8"	Fatty tissue over anterolateral thigh muscle
Children 12 mos or older, adolescents, and adults	5/8"	Fatty tissue over anterolateral thigh muscle or fatty tissue over triceps
Intramuscular (IM) injection Use a 22–25 gauge needle. Choose the injection site and needle length that is appropriate to the person's age and body mass.		
AGE	NEEDLE LENGTH	INJECTION SITE
Newborns (1st 28 days)	5/8"	Anterolateral thigh muscle
Infants (1–12 mos)	1"	Anterolateral thigh muscle
Toddlers (1–2 years)	1–1 1/4" 5/8–1"	Anterolateral thigh muscle or deltoid muscle of arm
Children and teens (3–18 years)	5/8–1"* 1–1 1/4"	Deltoid muscle of arm or anterolateral thigh muscle
Adults 19 years or older		
Female or male <130 lbs	5/8–1"*	Deltoid muscle of arm
Female or male 130–152 lbs	1"	Deltoid muscle of arm
Female 153–200 lbs Male 130–260 lbs	1–1 1/2"	Deltoid muscle of arm
Female 200+ lbs Male 260+ lbs	1 1/2"	Deltoid muscle of arm


* A 5/8" needle may be used for patients weighing less than 130 lbs (<60 kg) for IM injection in the deltoid muscle **only** if the skin stretched tight, the subcutaneous tissue is not bunched, and the injection is made at a 90-degree angle.

NOTE: Always refer to the package insert included with each biologic for complete vaccine administration information. CDC's Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well. Access the ACIP recommendations at www.immunize.org/acip.


Intramuscular (IM) injection



Subcutaneous (Subcut) injection



Intradermal (ID) administration of Fluzone ID vaccine



Intranasal (NAS) administration of Flumist (LAIV) vaccine

